



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/00, 31/55, 31/335, 31/215 // (A61K 31/55, 31:215, 31:17, 31:165) (A61K 31/55, 31:335, 31:17, 31:165)	A1	(11) International Publication Number: WO 98/20867 (43) International Publication Date: 22 May 1998 (22.05.98)
(21) International Application Number: PCT/US97/21723 (22) International Filing Date: 14 November 1997 (14.11.97) (30) Priority Data: 60/030,999 15 November 1996 (15.11.96) US (71) Applicant (for all designated States except US): THE UNITED STATES OF AMERICA, represented by THE SECRETARY, DEPARTMENT OF HEALTH AND HUMAN SERVICES National Institutes of Health [US/US]; Office of Technology Transfer, Suite 325, 6011 Executive Boulevard, Rockville, MD 20852 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): BLUMBERG, Peter, M. [US/US]; 4723-C Reels Mill Road, Frederick, MD 21704-7324 (US). BIRO, Tamas [HU/US]; 4740 Bradley Boulevard #319, Chevy Chase, MD 20815 (US). ACS, Peter [HU/US]; 4740 Bradley Boulevard #203, Chevy Chase, MD 20815 (US). ACS, Geza [HU/HU]; Pechy M.U. 28, H-4032 Debrecen (HU).		(74) Agents: GAGALA, Bruce, M. et al.; Leydig, Voit & Mayer, Ltd., Suite 4900, Two Prudential Plaza, 180 North Stetson, Chicago, IL 60601-6780 (US). (81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: PHARMACEUTICAL COMPOSITIONS CONTAINING VANILLOID AGONISTS IN COMBINATION WITH VANILLOID ANTAGONISTS (57) Abstract <p>The invention relates to pharmaceutical combinations containing capsaicin agonists with capsaicin antagonists for desensitization of vanilloid-responsive cells (a delta) and c-fiber sensory afferent neurons, and for increasing the therapeutic index of capsaicin agonists. Said combinations may contain for example compounds of formula (II) such as tinyatoxin or resiniferatoxin or compounds of formula (I) (as capsaicin agonists) together with capsazepin or other capsaicin antagonists of formula (III).</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

TECHNICAL FIELD OF THE INVENTION

10 BACKGROUND OF THE INVENTION

Resiniferatoxin (RTX), tinyatoxin (TTX), and related pharmacologically active compounds are potent stimulants of capsaicin-sensitive neurons. RTX, TTX, and analogues thereof are known in the art and described in U.S. Patent 4,939,149 (Blumberg) and in U.S. Patent 5,021,450 (Blumberg), both hereby
20 incorporated in their entirety by reference. RTX, TTX, and analogues thereof described above and below, stimulate and then desensitize vanilloid-sensitive neurons, in a manner similar to that of capsaicin. Therefore RTX, TTX, and their analogues have medical and veterinary utility in the treatment of arthritis, asthma, bladder hyperreflexia, allergic responses including rhinitis, fever, pain (including
25 pain associated with cancer, peripheral neuropathies, and postherpetic neuralgia), and in biological processes mediated by tachykinins, including substance P and other neuropeptides depleted by capsaicin.

1

A wide-variety of laboratory tests have also been used to demonstrate sensitivity to and potency of capsaicin-like molecules in the laboratory. These tests include the tail-flick test, yeast-induced inflammation test, the ethanol-induced gastric lesion test (Reid, supra). More specific tests employing ³H-resiniferatoxin binding to membrane preparations or cultured dorsal root ganglions (G. Acs et al. *Molecular Brain Research*, 35, 173-182 (1996)), as well as ion flux studies and electrophysiological studies as described by Bevan et al. (Bevan et al., *British Journal of Pharmacology*, 107, 544-552 (1992), hereby incorporated by reference) are also known.

Animals which exhibit a physiological or biochemical sensitivity to capsaicin or RTX are referred to as capsaicin-sensitive. Additionally neurons and other cells that bind or respond to capsaicin, or RTX, are referred to as capsaicin-sensitive neurons. However, capsaicin-sensitive neurons can be defined more specifically as those neurons that not only respond to capsaicin, but also become insensitive (desensitized) to capsaicin or RTX upon repeated capsaicin challenge (Buck et al., *Pharmacol. Rev.*, 38, 179-226 (1986) and Holzer, *Pharmacol. Rev.*, 43, 144-201 (1991)). It is this latter, more specific definition which is preferred herein.

It is generally understood in the art that the homovanillic acid moiety shared by capsaicin and RTX is central to the similarity of the pharmacological activity of these neuroactive compounds. Accordingly, the site of capsaicin activity is commonly referred to as a vanilloid receptor.

Capsaicin-like antagonists of capsaicin are also known. The most familiar example of these antagonists is capsazepine. Capsazepine and molecules similar to capsazepine are described in Walpole et al., *J. Med. Chem.*, 37, 1942-1954 (1994), hereby incorporated by reference in its entirety. Capsazepine and related compounds are competitive antagonists of both capsaicin and capsaicin-like agonists and RTX and RTX-like agonists.

Despite the known properties of capsaicin and RTX, and the known properties of capsaicin-like antagonists, there has been no motivation heretofore to use a therapeutic combination of RTX-like capsaicin agonists and capsaicin-like antagonists to desensitize a vanilloid-responsive cell. In fact, heretofore it was believed that the acute response to capsaicin agonists was causally linked to the desensitization that follows. Prior to the present disclosure it would have been believed that administration of antagonists in sufficient quantities to attenuate the toxic response to capsaicin agonists would have attenuated the subsequent desensitization to an equal extent.


Accordingly it is a principle object of the present invention to provide a pharmaceutical composition which includes a therapeutically effective combination of RTX-like capsaicin agonists and capsaicin-like antagonists that is capable of desensitizing a vanilloid responsive cell with a reduced toxic response normally attendant the administration of the agonist alone.

It is another object of the present invention to provide a method for desensitizing a vanilloid-responsive cell which includes the administration of a therapeutically effective combination of a RTX-like capsaicin agonist and a capsaicin-like antagonist.


It is yet another object of the invention to provide a pharmaceutical composition as described which has attenuated side effects. These and other objects and advantages of the present invention will be apparent from the following detailed description of the invention.

SUMMARY OF THE INVENTION

The present invention is predicated on the discovery that vanilloid-responsive cells can be desensitized by a therapeutically effective combination of a capsaicin agonist and a capsaicin antagonist. Advantageously, the therapeutic index of the combination therapy exceeds the therapeutic index of administering the agonist alone. The present invention thus provides a method of desensitizing a vanilloid-responsive cell comprising administering to a capsaicin-sensitive animal a therapeutically effective combination of capsaicin agonist and a capsaicin antagonist. In accordance with the invention, the capsaicin agonist can be of the formula




Formula I




- wherein R_9 is hydroxy or methoxy, R_{10} and R_{11} , which may be the same or different, are independently selected from the group consisting of hydrogen, hydroxyl, or a C_{1-10} alkyl or aryl ester, R_{12} , R_{14} , and R_{15} , which may be the same or different, are independently selected from the group consisting of hydrogen, hydroxy, methoxy, sulfhydryl, nitro, amino, ethoxy, halo, $OCOCH_3$, and $O(CH_2)_2NH_2$, R_{13} is selected from the group consisting of hydroxy, $OCOCH_3$, and $O(CH_2)_2NH_2$.

The capsaicin agonist can also be of the formula



Formula II






- wherein R_2 is hydroxy or methoxy, R_3 is a loweralkylaryl or aryl group having 1 to 3 rings, R_4 to R_7 , which may be the same or different, are independently selected from the group consisting of hydrogen, hydroxy, methoxy, sulfhydryl, nitro, amino, ethoxy, halo, and OCOCH_3 .

In accordance with the invention, the antagonist of capsaicin is of the formula



wherein:





and wherein J_1 is independently selected from the group consisting of C_{1-3} alkoxy, hydroxy, a halo, and a divalent alkyl which together with a nitrogen of X forms a heterocyclic ring of from 5 to 12 carbon atoms and wherein, when n is greater than one, each J_1 may be the same or different; n is 1-5; J_2 and J_3 are independently selected from hydrogen, and C_{1-5} alkyl; D is selected from the group consisting of C_{4-20} alkyl, C_{4-20} alkenyl, C_{4-20} alkynyl, aryl of 1-3 rings substituted with C_{1-8} alkyl, a C_{4-20} alkyl, alkenyl or alkynyl substituted with an aryl of 1 to 3 rings, aryl, haloaryl, haloarylalkenyl, such that the vanilloid-responsive cell becomes less sensitive to an initial administration of capsaicin and such that the acute response to a capsaicin agonist is attenuated.

The present invention also provides a pharmaceutical composition comprising the an agonist of Formula I or Formula II, an antagonist of Formula III, and a pharmaceutically acceptable carrier.

It will be appreciated that the capsaicin agonist and the antagonist can be used concurrently, e.g., in the same pharmaceutical composition or administered at the same time, or the agonist and antagonist may be used sequentially to reduce the toxic effects of the agonist and/or increase the therapeutic index.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 depicts the data demonstrating the enhanced therapeutic index of resiniferatoxin when capsazepine is co-administered.

DETAILED DESCRIPTION OF THE INVENTION

As observed for capsaicin, treatment of vanilloid sensitive cells with resiniferatoxin, and related compounds, causes an initial excitation and a subsequent desensitization to repeated applications of the compound (Szallasi et al, Neuroscience, 30, 515-520 (1989), Blumberg et al., Resiniferatoxin, An Ultrapotent Capsaicin Analog, in Capsaicin in the Study of Pain, J.N. Wood, ed., pages 45-62, Academic Press Publishers, London (1993)). The initial excitation is frequently manifested as pain in the patient or subject and can be quantified by

directly or indirectly measuring the calcium ion flux induced in vanilloid sensitive cells. Desensitization is a blunting or abrogation of response to capsaicin agonists measured at a time point a few hours after than the initial application of the agonist (e.g., 2-8 hours later, six hours later is a common time point for conducting this measurement). For example, desensitization can be effected in some cells with 250 pM RTX. Subsequent dosages of capsaicin agonists nearly 1000 times more potent (e.g., 200 nM RTX) will then fail to induce a strong response in vanilloid sensitive cells. This is a typical illustration of desensitization, and is believed to be the mechanism by which the commercially available analgesic Zostrix™ (capsaicin) functions.

It has been postulated for decades that the pungency of capsaicin analogs is proportional to the desensitization that follows (Jansco, Desensitization With Capsaicin and Related Acylamides as a Tool for Studying the Function of Pain Receptors, In Pharmacology of Pain, 9, 33-55, Pergamon Publishers, Oxford, U.K., (1968)). However, quite unexpectedly and surprisingly it has been found that the pungency of resiniferatoxin, tinyatoxin, related daphnanes, and C-20 homovanillyl phorbolesters is not proportional to the desensitization that follows.


It has now been found that resiniferatoxin, tinyatoxin, and related daphnanes and C-20 homovanillyl phorbol esters induce desensitization to further administration of capsaicin agonists, or other painful stimuli by a process which is distinct from the acute response (i.e., the initial excitation, sometimes referred to as the toxic response) initially induced by these compounds. Further, it has now been found that RTX, and the like, are capable of inducing desensitization even in the presence of competing concentrations of capsazepine and other antagonists, which are capable of blocking or significantly attenuating the acute phase response to the drug. Accordingly, in accordance with the invention it is possible to formulate pharmaceutical compositions with an enhanced therapeutic index for clinical and veterinary use in the desensitization of vanilloid sensitive cells by controlling the proportion of an RTX-like agonist with a capsaicin-like antagonist.

As used herein, the therapeutic index of a pharmaceutical composition is a number representing the magnitude of desensitization produced by the administration of the pharmaceutical composition divided by the magnitude of the acute (toxic) response induced by the administration of the pharmaceutical composition. It is often useful to substitute the level of calcium ion influx in capsaicin responsive neurons for the direct measurement of response to resiniferatoxin (or the like) during the initial application of such capsaicin agonists, because calcium ion influx is more easily quantified than pain and other


indications of acute phase response. Similarly, desensitization can be quantified by observing the degree of inhibition of calcium ion flux upon a subsequent challenging application of capsaicin or the like.

5 The pharmaceutical compositions can be prepared as one discrete formulation or can be formulated from two or more discrete preparations. If the pharmaceutical composition is composed of multiple components, these components can be mixed before administration to an animal or may be co-administered to achieve the desired therapeutic benefit. Regardless of the initial form of the pharmaceutical composition, the pharmaceutical composition is
10 formulated and administered such that the animal receives a capsaicin agonist and a capsaicin antagonist dissolved, suspended, or dispersed in a pharmaceutically acceptable carrier.


15 The capsaicin agonist which mediates the therapeutic desensitization is of the formula




Formula I



- wherein R_9 is hydroxy or methoxy, R_{10} and R_{11} , which may be the same or different, are independently selected from the group consisting of hydrogen, hydroxyl, or a C_{1-10} alkyl or aryl ester (such as decanoate, benzoate, or phenylacetate), R_{12} , R_{14} , and R_{15} , which may be the same or different, and are independently selected from the group consisting of hydrogen, hydroxy, methoxy, sulfhydryl, nitro, amino, ethoxy, halo (i.e., fluoro, chloro, bromo, iodo), $OCOCH_3$, and $O(CH_2)_2NH_2$, R_{13} is selected from the group consisting of hydroxy, $OCOCH_3$, and $O(CH_2)_2NH_2$, or is of the formula



Formula II



- wherein R_2 is hydroxy or methoxy, R_3 is a loweralkylaryl or aryl group having
 5 from 1 to 3 rings, preferably an aryl group with 1 ring having 5-8 ring members,
 yet more preferably phenyl or C_{1-4} alkylphenyl, R_4 to R_7 , may be the same or
 different, and are independently selected from the group consisting of hydrogen,
 hydroxy, methoxy, sulfhydryl, nitro, amino, ethoxy, halo (i.e., chloro, fluoro,
 bromo, iodo), $OCOCH_3$. By loweralkyl is meant a alkyl moiety of 1 to 8 carbons,
 10 preferably 1 to 4 carbons.


- One particular example of a compound of Formula II is tinyatoxin.
 Another particular example of a compound of Formula II is resiniferatoxin. Use of
 resiniferatoxin is advantageous in that it is among the capsaicin analogs most
 15 widely utilized in current biomedical research. This research has revealed that
 resiniferatoxin is up to 20,000 times more potent than capsaicin. Surprisingly, it
 has now been found that competitive antagonists of resiniferatoxin and capsaicin
 (e.g., capsazepine) can inhibit the toxic effects of resiniferatoxin administration at
 least about ten-fold more effectively than they inhibit the desensitizing effects of
 20 resiniferatoxin. Therefore, the combination of resiniferatoxin, or the like, with
 capsazepine or the like, allows a greater therapeutic benefit to be derived from the
 administration of resiniferatoxin, because it is the toxic effects of capsaicin and
 resiniferatoxin that function to limit the amount of capsaicin or resiniferatoxin
 which can be administered to a patient and these toxic effects are suitably reduced

when capsazepine-like antagonists are co-administered with resiniferatoxin or the like.



The capsaicin antagonists useful accordance with the present invention are
5 of the formula



wherein:



10



15 and wherein J_1 is independently selected from the group consisting of C_{1-3} alkoxy, hydroxy, a halo, and a divalent alkyl which together with a nitrogen of X forms a heterocyclic ring of from 5 to 12 carbon atoms and wherein when n is greater than one, each J_1 may be the same or different; n is 1-5; J_2 and J_3 are independently
20 selected from hydrogen, and C_{1-5} alkyl; D is selected from the group consisting of C_{4-20} alkyl, C_{4-20} alkenyl, C_{4-20} alkynyl, aryl of 1-3 rings substituted with C_{1-8} alkyl, a C_{4-20} alkyl, alkenyl or alkynyl substituted with an aryl of 1 to 3 rings, aryl, haloaryl, haloarylalkenyl, such that the vanilloid-responsive cell becomes less sensitive to an initial administration of capsaicin and such that the acute response
25 to a capsaicin agonist is attenuated.

Compounds illustrative of Formula III include the hydroxy-halo benzene containing compounds, which include compounds of Formula III wherein n is 2 or 3, preferably 2, and a first occurrence of J_1 is hydroxy, and a second occurrence of

J₁ is halo, preferably chloro. Commonly, the hydroxy moiety is 4-hydroxy, and the halogen, e.g., is 3-chloro.


Another group of compounds illustrative of Formula III include methoxybenzene containing compounds of Formula III, wherein n is 1 or 2, preferably 1, and at least one occurrence of J₁ is methoxy, preferably 3-methoxy.

Another group of compounds illustrative of Formula III include the divalent-alkylbenzene containing compounds of Formula III. A well known example of this group is capsazepine. This group of compounds are illustrated by compounds wherein a first occurrence of J₁ is 2, N-propyl and additional occurrences of J₁ include 3-hydroxy and 4-hydroxy. A particularly preferred characteristic of the divalent-alkylbenzene containing compounds of Formula III are those in which the aryl ring of A is perpendicular the plane of the amide or thiourea of B. By way of illustration, the A ring is usually perpendicular to the plane of B when the ring formed in part by the divalent alkyl moiety has seven ring atoms and is attached at position 2 of the A ring.


Other compounds of Formula III of particular interest are those in which J₂ is hydrogen.

Moreover, while the hydrophobic moiety represented by D in Formula III can vary, those D moieties of particular interest are a C₆₋₁₀ alkyl, alkenyl, or alkynyl, especially about C₈ alkyl, and substituted or unsubstituted aryl alkenyl moieties such as iodophenylpropenamide.


Particular embodiments of Formula III that are useful in the context of the present invention include N-octyl N'(p-hydroxy m-chlorobenzyl) thiourea, N-octyl N'(p-hydroxy m-chlorophenyl)acetamide, N-(o-methoxybenzyl)-N'-octyl thiourea, and 3-methoxybenzyl p-iodophenylpropenamide. The best known antagonist suitable for use in the present invention is capsazepine. To better illustrate the compounds of Formula III the following depictions are provided.



3-methoxybenzyl p-iodophenylpropenamide




N-(o-methoxybenzyl), N'-octyl thiourea




N-octyl (p-hydroxy m-chlorophenyl) acetamide

5



N-octyl, N'-(p-hydroxy m-chlorobenzyl) thiourea



Capsazepine

10

Advantageously, any one or more compounds of Formula I or Formula II can be used in combination to practice the claimed method of desensitizing an vanilloid-responsive cell and, any one or more compounds of Formula I or

15 Formula II can be combined with any one or more compound of Formula III in a pharmaceutically acceptable carrier.

The compounds useful in the context of the present inventive method can be administered to a capsaicin-sensitive animal, especially humans, non-human primates, horses, dogs, rats, and mice by any suitable means. Such administration means or routes include for example, orally, inhalation, parenterally, vaginally, rectally, intranasal, intravesicular, intraperitoneally, and, especially, topically.

When applied topically the agonist should be present in a concentration sufficient to provide a therapeutic benefit (i.e., desensitization) in the presence of the antagonist. In one particular embodiment, this amount is from a 0.0001% to a 1% cream, preferably a 0.025% to 0.1% cream. To effect a suitable attenuation of the initial excitement without abrogating the desensitization, the relative strength of the antagonist (e.g., capsazepine) must be balanced against the relative strength of the antagonist. For example, if a relatively very weak agonist is selected for use with a strong antagonist, then the ratio of antagonist to agonist will be smaller than if a relatively strong agonist is used with the same antagonist. Accordingly, ratios of antagonist to agonist varying in the range of 1:1 up to 10^8 :1 are contemplated. Advantageously, standard potencies can be derived by comparison to capsaicin or resiniferatoxin for agonists and by comparison to capsazepine for antagonists. After making such a routine determination, calculating the appropriate ratios for use in the context of the present invention would be straightforward. Specific ratios for using resiniferatoxin in combination with capsazepine and the rationale for selecting these ratios are disclosed in detail below.

When administered internally, the quantity of agonist and antagonist will vary according to the distribution properties associated with the pharmaceutically acceptable carrier or carriers and the distribution kinetics associated with that carrier when administered by a particular route in the patient or subject. The ratio of agonist to antagonist, however, should remain essentially constant. If the internally applied composition will be dispersed in the blood stream (e.g., if applied in liposomes administered intravenously) then one skilled in the art will appreciate that 6-8% of the weight of the patient will be derived from the blood volume. Accordingly, an agonist with a particular potency should be administered such that it is equivalent in its agonism to about 25 pM to about 25 nM resiniferatoxin and an antagonist with a particular potency should be administered such that it is equivalent in its antagonism to between 1 μ M to 1 mM capsazepine. In one particular embodiment, this value will be approximately 10 to 35 times the dose required for maximal inhibition of acute phase response to capsaicin. Of course, the amount administered will vary according to such parameters as the severity of the indication treated, the age, weight, and general health of the patient

or subject, the carrier employed, and the route of administration. Further, the quantity employed will be in keeping with the entire therapeutic regime for an animal under the care of a clinician or veterinarian.

5 A therapeutically effective amount of the combined pharmacologically active ingredients of the present invention can be used in a pharmaceutical composition, which of course, will depend upon the particular means or route of administration, as well as other practical considerations. Accordingly, there is a wide variety of suitable formulations of the pharmaceutical compositions of the present invention.

10 Formulations suitable for oral administration can comprise (a) liquid solutions, such as water or saline, (b) capsules, sachets or tablets, each containing a predetermined amount of the active ingredient, as solids or granules, (c) suspensions in an appropriate liquid, and (d) suitable emulsions. Tablet forms can include one or more of lactose, mannitol, corn starch, potato starch,
15 microcrystalline cellulose, acacia, gelatin, colloidal silicon dioxide, croscarmellose sodium, talc, magnesium stearate, stearic acid, and other excipients, colorants, diluents, buffering agents, moistening agents, preservatives, flavoring agents, and pharmacologically compatible carriers. Lozenge forms can comprise the active ingredient in a flavor, usually sucrose and acacia or tragacanth, as well as pastilles
20 comprising the active ingredient in an inert base, such as gelatin and glycerin or sucrose and acacia emulsions, gels, and the like containing, in addition to the active ingredient, such carriers as are known in the art.

One particular embodiment among the preferred embodiments of the present invention comprises a combination of a resiniferatoxin-like capsaicin
25 agonist, in combination with a capsazepine-like capsaicin antagonist, in a cream, paste, emulsion, or the like for application to the skin of a human, domesticated horse, or a dog kept as a pet (as opposed to a research animal), especially to the skin of a human. The dosages of the pharmacologically active ingredients for such a pharmaceutical composition are disclosed above.

30 All pharmaceutical compositions disclosed above can comprise formulations may include diluents, such as water and alcohols, for example, ethanol, benzyl alcohol, and the polyethylene alcohols, either with or without the addition of a pharmaceutically acceptable surfactant, suspending agent, or emulsifying agent. Capsule forms can be of the ordinary hard- or soft-shelled
35 gelatin type containing, for example, surfactants, lubricants, and inert fillers, such as lactose, sucrose, calcium phosphate, and corn starch. Tablet forms can include one or more of lactose, sucrose, mannitol, corn starch, potato starch, alginic acid,

microcrystalline cellulose, acacia, gelatin, guar gum, colloidal silicon dioxide, croscarmellose sodium, talc, magnesium stearate, calcium stearate, zinc stearate, stearic acid, and other excipients, colorants, diluents, buffering agents,

5 disintegrating agents, moistening agents, preservatives, flavoring agents, and pharmacologically compatible excipients. Lozenge forms can comprise the active ingredient in a flavor, usually sucrose and acacia or tragacanth, as well as pastilles comprising the active ingredient in an inert base, such as gelatin and glycerin, or sucrose and acacia, emulsions, gels, and the like containing, in addition to the active ingredient, such excipients as are known in the art.

10 Regardless of the form of the combined capsaicin agonists and antagonists, the present invention provides a method of desensitizing a vanilloid-responsive cells that employs administering to a capsaicin-sensitive animal a therapeutically effective combination of capsaicin agonist and an antagonist of capsaicin such that the vanilloid-responsive cell becomes less sensitive to further or repeated
15 administration of capsaicin and such that the acute response to the capsaicin agonist is attenuated.

Many but not all animals are sensitive to vanilloids. Humans, monkeys, horses, dogs, rats, and mice are all suitably sensitive to vanilloids. It is routine to determine whether any other animal is also sensitive to vanilloids such as capsaicin
20 and resiniferatoxin. Ingestion of vanilloids in sensitive animals often causes gustatory sweating, while application to mucosa or skin is often accompanied by a burning sensation or behavior which would indicate the perception of a burning sensation (e.g., tail-flicking or eye-wiping).

Cells which are sensitive to vanilloids as meant herein exhibit a calcium ion
25 influx upon a first exposure to capsaicin and the like upon a first administration and exhibit a reduced response to similar or even elevated concentrations of capsaicin a short period later (e.g., 2 to 8 hours later, often measured 6 hours later). The calcium uptake can be measured by any suitable means. Suitable measurements include behavioral measurements such as eye-wiping,
30 electrophysiological measurement of calcium-dependent ion flows, or monitoring of the uptake of radioactive calcium.

Primates are frequently about ten-fold less sensitive to vanilloids than rats. Therefore, optimum concentrations of either the agonists or antagonists for primates, including humans will typically relate to optimum concentrations
35 observed for rats by 0.1 to 1000 times, preferably 5 to 100 times, and most preferably 5 to 50 times the concentrations.

Particular examples of vanilloid responsive cells in rats and humans are sensory afferent neurons, such as C-fiber sensory afferent neurons or a A δ -fiber sensory afferent neuron. It is also known in the art that some neurons of the central nervous system contain vanilloid receptors and can be treated in accordance with the present invention.

In selecting a dosage of capsaicin antagonist, it is often useful to select a dosage that is 10-30 times the dose that is required to inhibit an acute response to capsaicin. In this regard, it is useful to determine either the dosage of antagonist at which less than 10% of the maximum calcium ion influx occurs in capsaicin sensitive cells and multiply that dosage by between 10 and 30 times or to determine a dosage 10-30 times the K_i with respect to an optimal dose of capsaicin. Since the antagonist capsazepine has been studied more extensively than any other capsaicin antagonist, its use as the antagonist in the present inventive composition and method is among the preferred embodiments.

Example 1

The following example demonstrates the relative affinity of various compounds useful in the context of the present invention for inhibiting the specific binding of ³H-resiniferatoxin to rat dorsal root ganglion membrane preparations and for stimulation of calcium (⁴⁵Ca) uptake in cultured rat dorsal root ganglion neurons.

Female Sprague-Dawley rats (150-160 g body weight) were allowed access to food and water ad libitum. Anesthetized rats were killed by decapitation. The spinal columns were removed aseptically and dorsal root ganglia were dissected out and collected in DMEM containing 0.5% heat inactivated fetal bovine serum, 25 mM HEPES, 1 mM Na-pyruvate, and antibiotics. Ganglia were digested with 0.125% collagenase solution, washed twice in medium and triturated to form a single cell suspension. The cells were pelleted through a cushion of DMEM containing 15% fatty acid free BSA to remove myelin debris.

Resuspended cells were plated in laminin-coated 24-well plates at a density of 2 x 10⁴ cells/well in 500 μ l medium, treated with 10⁻⁴ M cytosine arabinoside for 24 hours, and cultured for 5 days in DMEM containing 10% fetal bovine serum, 10% horse serum, antibiotics, and 200 ng/ml mouse submaxillary gland 2.5S nerve growth factor.

Alternatively, dispersed ganglia from six rats were placed in two 150 mm laminin coated dishes to achieve neuronal enrichment. After 18 hours, the non-

neuronal cells were firmly attached to the dish. Neuronal cells, which were weakly adherent, were selectively dislodged, washed in DMEM and re-plated at a density of 5×10^3 cells/well (96 well plate) in 100 μ l serum-free medium and used for resiniferatoxin binding assays and calcium uptake assays immediately.

- 5 Membrane preparations were then prepared as described by Szallasi et al., Resiniferatoxin In: P.M. Conn (Ed.), Methods in Neurosciences, 8, 368-380 Neurotoxins, Academic Press, Orlando, FL, USA, (1990). Tritiated resiniferatoxin binding assays were carried out as described, *ibid.* Six to 500 pM ^3H -resiniferatoxin was used for binding analysis. Nonspecific binding was
10 determined in the presence of 100 nM radioinert resiniferatoxin. Binding data used to calculate the data of Table 1 were expressed as fmol/mg protein.

- Calcium uptake by these ganglia was determined by culturing the ganglia as above and treating with 1 $\mu\text{Ci/ml}$ ^{45}Ca and increasing concentrations of the different compounds. Cells were washed six times by filtration. Calcium uptake
15 was determined by scintillation counting.

Table 1. Effectiveness of various vanilloids in rat dorsal root ganglion cells.

	Inhibition of ^3H -RTX binding		Stimulation of Calcium Uptake	
	Hill		Hill	
	K_i (nM)	Coefficient	EC_{50}	Coefficient
RTX	0.0385	1.95	0.94	1.15
RTX-amide ¹	25.7	1.92	26.5	1.07
RTX-thiourea ²	68.5	1.98	149	0.93
TTX	0.173	1.82	1.46	1.18
PDB-HVA ³	7980	1.72	1410	1.11
Capsaicin	4930	1.81	340	1.08
Capsazepine	3890	1.74	~300	0.96

- 20
1 Resiniferonol 20-homovanillylamide
2 4-hydroxy-3-methoxyphenyl thiourea
3 Phorbol 12,13-dibenzoate 20-homovanillylamide

Example 2

- 25 The following example compares the sensitivity of rats and monkeys to resiniferatoxin mediated desensitization, resiniferatoxin mediated calcium uptake, and capsaicin mediated calcium uptake. This information allows one skilled in the art to extend the data derived from rats to the treatment of primates, especially monkeys and humans.

- 30 Using methods similar to those described in Example 1, the data of Table 2 were collected. These data show that monkeys are also capsaicin-sensitive, but are approximately 3-6 fold less sensitive than rats. Therefore, the present inventive

pharmaceutical compositions should be formulated 3-6 fold more concentrated with respect to the active ingredients when used for monkeys, rather than rats.

Table 2. Relative sensitivity to vanilloids of rat and monkey.

5

	<u>Rat</u>	<u>Monkey</u>	<u>Ratio</u>
RTX desensitization	0.081 nM	0.46 nM	5.7
RTX Ca uptake	1.24 nM	6.2 nM	5.0
Capsaicin Ca uptake	316 nM	880 nM	2.8

Example 3

This example sets forth the effect of pre-treating vanilloid sensitive cells with resiniferatoxin.

10

Ganglia, prepared as in Example 1, were subjected to varying concentrations of resiniferatoxin either six hours preceding administration of 3 μ M capsaicin or simultaneously with 3 μ M capsaicin. No utilized concentration of resiniferatoxin had any effect on calcium uptake in response to 3 μ M capsaicin when the compounds were co-administered. When cells were pretreated with

15

concentrations of resiniferatoxin ranging from 0.1 to 0.5 nM, a dose dependent inhibition of calcium uptake was observed upon subsequent challenge with 3 μ M capsaicin (six hours later). Treatments utilizing between 0.5 to 2.0 nM resiniferatoxin, were sufficient to completely block detectable calcium uptake upon challenge with capsaicin six hours later.

20

Example 4

This example sets forth that co-administration of a capsaicin antagonist (e.g., capsazepine) with a capsaicin agonist (e.g., resiniferatoxin) results in effective desensitization of vanilloid sensitive cells with a concomitant reduction

25

in acute phase toxicity. That is, co-administration of antagonists with agonists results in an increase in the therapeutic index relative to the administration of agonists alone.

30

Control and capsazepine pretreated cells were challenged with different concentrations of resiniferatoxin to induce 45 Ca uptake. The data points derived from these experiments are depicted in Figure 1 by squares. It is apparent that the co-administration of capsazepine (indicated by filled squares) increases the amount of resiniferatoxin required to induce a given level of calcium uptake by about ten-fold relative to cells not administered capsazepine (indicated by open squares. The large shift to the right of the data from the cells treated with capsazepine indicates

that there is a reduction in the calcium influx mediated by moderate levels of resiniferatoxin in the acute (toxic) phase of administration. That is, e.g., a rat administered 3 nM resiniferatoxin would experience severe pain in the absence of 10 μ M capsazepine, but nearly no pain in the presence of the capsazepine.

5 The effect on desensitization of co-administration of capsazepine was also examined. In Figure 1, the circles indicate the reduction in the level of calcium uptake as a function of pretreatment with a given concentration of resiniferatoxin and when challenged six hours later with 3 μ M capsaicin. The filled circles depict the results when the cells are co-administered 10 μ M capsazepine and the
10 indicated concentration of resiniferatoxin, and the open circles depict the results from when the cells are administered resiniferatoxin alone. The small degree of a shift to the right of the line formed by the open circles relative to the line formed by the filled circles indicates that the (therapeutic) desensitization mediated by pretreatment with 100 to 250 pM resiniferatoxin is not adversely affected by the
15 co-administration of capsazepine.



 All the data depicted in Figure 1, therefore show that the presence of capsazepine increases the therapeutic index of resiniferatoxin by about 10-fold. These data also indicate that a ratio of 100 to 250 pM resiniferatoxin to about 10 μ M capsazepine effectively increases the therapeutic index of resiniferatoxin. It
20 should be appreciated that this ratio is central to the functionality of the instant embodiment of the present invention, rather than the particular quantity of resiniferatoxin or capsazepine. While not being bound by any particular theory, this appears to be true because of the competitive nature of capsazepine inhibition.

25 All of the references cited herein, including patents, patent applications, and publications, are hereby incorporated in their entireties by reference.

 While this invention has been described with an emphasis upon preferred embodiments, it will be obvious to those of ordinary skill in the art that variations of the preferred embodiments may be used and that it is intended that the invention
30 may be practiced otherwise than as specifically described herein. Accordingly, this invention includes all modifications encompassed within the spirit and scope of the invention as defined by the following claims.

What is claimed is:


1. A pharmaceutical composition comprising a capsaicin agonist, a capsaicin antagonist, and a pharmaceutically acceptable carrier.
2. The pharmaceutical composition of claim 1, wherein said agonist is of the formula






wherein R_9 is hydroxy or methoxy, R_{10} and R_{11} , which may be the same or different, are independently selected from the group consisting of hydrogen, hydroxyl, or a C_{1-10} alkyl or aryl ester, R_{12} , R_{14} , and R_{15} , which may be the same or different, are independently selected from the group consisting of hydrogen, hydroxy, methoxy, sulfhydryl, nitro, amino, ethoxy, halo, $OCOCH_3$, and

$\text{O}(\text{CH}_2)_2\text{NH}_2$, R_{13} is selected from the group consisting of hydroxy, OCOCH_3 , and $\text{O}(\text{CH}_2)_2\text{NH}_2$.

3. The pharmaceutical composition of claim 1, wherein said agonist is of the formula



Formula II




wherein R_2 is hydroxy or methoxy, R_3 is a loweralkylaryl or aryl group having from 1 to 3 rings, R_4 to R_7 are independently selected from the group consisting of hydrogen, hydroxy, methoxy, sulfhydryl, nitro, amino, ethoxy, halo, OCOCH_3 .

4. The pharmaceutical composition of claim 3, wherein said agonist is tinyatoxin.




5. The pharmaceutical composition of claim 3, wherein said agonist is resiniferatoxin.

6. The pharmaceutical composition of any of claims 1-5, wherein said antagonist is of the formula



Formula III

wherein:



and wherein J_1 is independently selected from the group consisting of C_{1-3} alkoxy, hydroxy, a halo, and a divalent alkyl which together with a nitrogen of X forms a heterocyclic ring of from 5 to 12 carbon atoms and when n is greater than one, each J_1 may be the same or different; n is 1-5; J_2 and J_3 are independently selected from the group consisting of hydrogen and C_{1-5} alkyl, D is selected from the group consisting of C_{4-20} alkyl, C_{4-20} alkenyl, C_{4-20} alkynyl, aryl of 1-3 rings substituted with C_{1-8} alkyl, a C_{4-20} alkyl, alkenyl or alkynyl substituted with an aryl of 1 to 3 rings, aryl, haloaryl, and haloarylalkenyl.

7. The pharmaceutical composition of claim 6, wherein said antagonist is selected from the group consisting of N-octyl N'(p-hydroxy m-chlorobenzyl) thiourea, N-octyl N'(p-hydroxy m-chlorophenyl)acetamide, N-(o-methoxybenzyl)-N'-octyl thiourea, and 3-methoxybenzyl p-iodophenylpropenamide.

8. The pharmaceutical composition of claim 6, wherein said antagonist is capsazepine.

9. A method of desensitizing a vanilloid-responsive cell comprising administering to a capsaicin-sensitive animal, said capsaicin agonist and said capsaicin antagonist of the pharmaceutical compositions of any of claims 1-8.


10. The method of claim 9, wherein said vanilloid responsive cell is a C-fiber sensory afferent neuron or a A δ -fiber sensory afferent neuron.

11. The method of claim 9 or 10, wherein the dosage of antagonist is 10-30 times the dose that is required to inhibit an acute response to capsaicin.

12. The method of claim 9 or 10, wherein the dosage of antagonist is 10-30 times the dosage required to result in a concentration approximating the K_i with respect to capsaicin.

13. A method of increasing the therapeutic index of a capsaicin agonist of Formula I or of Formula II comprising administering to a capsaicin sensitive animal a capsaicin antagonist of Formula III.

1/1

**FIG. 1**

INTERNATIONAL SEARCH REPORT

Intern. Appl. No.

PCT/US 97/21723

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K31/00 A61K31/55 A61K31/335 A61K31/215 //(A61K31/55,
31:215,31:17,31:165), (A61K31/55,31:335,31:17,31:165)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DICKENSON ET AL.: "Selective antagonism of capsaicin by capsazepine: evidence for a spinal receptor site in capsaicin-induced antinociception" BR. J. PHARMACOL., vol. 104, 1991, pages 1045-1049, XP002059189 see the whole document	1,6,8
Y	---	3,5,9-13
	-/--	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

17 March 1998

Date of mailing of the international search report

03.04.98

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Gac, G

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 97/21723

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DRAY ET AL.: "Antagonism of capsaicin-induced activation of C-fibres by a selective capsaicin antagonist, capsazepine" BR. J. PHARMACOL., vol. 99, 1991, page 78p XP002059190 see abstract	1,6,8
Y	---	3,5,9,10
X	BEVAN ET AL.: "Capsazepine: a competitive antagonist of the sensory neurone excitant capsaicin" BR. J. PHARMACOL., vol. 107, no. 2, October 1992, pages 544-552, XP002059191 see the whole document	1,6,8
Y	---	3,5,9,10
X	SZALLASI ET AL.: "Competitive inhibition by capsazepine of [3H] resiniferatoxin binding to central (spinal cord and dorsal root ganglia) and peripheral (urinary bladder and airways) vanilloid (capsaicin) receptors in the rat" J. PHARMACOL. EXP. THER., vol. 267, no. 2, 1993, pages 728-733, XP002059192 see the whole document	1,6,8
Y	---	3,5,9-13
X	SZOLCSANYI ET AL.: "Capsaicin-sensitive bronchopulmonary receptors with dual sensory efferent function: mode of action of capsaicin antagonists" ACTA PHYSIOL. HUNG., vol. 77, no. 3-4, 1991, pages 293-304, XP002059193 see the whole document	1,9,10
Y	---	1,3,5,6,8,13
A	WO 90 04389 A (UNITED STATES OF AMERICA) 3 May 1990 cited in the application see the whole document	9,10
P,Y	---	1,3,5,6,8,13
A	WO 97 17077 A (L'OREAL) 15 May 1997 see page 4 see page 5, line 4 - line 6 see page 10, line 12 - line 32 see page 11, line 16 - line 26	9

	-/--	

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 97/21723

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	ACS ET AL.: "Distinct structure-activity relations for stimulation of ⁴⁵ Ca uptake and for high affinity binding in cultured rat dorsal root ganglion neurons and dorsal root ganglion membranes" MOL. BRAIN RES., vol. 35, no. 1-2, 1996, pages 173-182, XP002059194 see the whole document	1-6,8-10
A	---	11,12
Y	SZALLASI ET AL.: "A novel agonist, phorbol 12-phenylacetate 13-acetate 20-homovanillate, abolishes positive cooperativity of binding by the vanilloid receptor" EUR. J. PHARMACOL., vol. 299, no. 1-3, 1996, pages 221-228, XP002059195 see the whole document	1-6, 8-10,13
Y	---	
Y	APPENDINO ET AL.: "Synthesis and evaluation of phorboid 20-homovanillates: discovery of a class of ligands binding to the vanilloid (capsaicin) receptor with different degrees of cooperativity" J. MED. CHEM., vol. 39, no. 16, 1996, pages 3123-3131, XP002059196 see the whole document	1-6, 8-10,13
A	---	
A	WALPOLE C S J ET AL: "THE DISCOVERY OF CAPSAZEPINE, THE FIRST COMPETITIVE ANTAGONIST OF THE SENSORY NEURON EXCITANTS CAPSAICIN AND RESINIFERATOXIN" JOURNAL OF MEDICINAL CHEMISTRY, vol. 37, no. 13, 24 June 1994, pages 1942-1954, XP000576031 cited in the application	1,3,5,6, 8
A	---	
A	WO 90 15055 A (THE UNITED STATES OF AMERICA) 13 December 1990 cited in the application see the whole document, especially page 11 lines 33-37 & US 5 021 450 A	1-10
A	---	
A	EP 0 068 590 A (THE PROCTER & GAMBLE COMPANY) 5 January 1983 see the whole document	6,7

	-/--	

1

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 97/21723

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>SZALLASI ET AL.: "Inhibition of [3H] resiniferatoxin binding to rat dorsal ganglion membranes as a novel approach in evaluating compounds with capsaicin-like activity"</p> <p>NAUNYN-SCHMIEDEBERG'S ARCH. PHARMACOL., vol. 344, no. 5, 1991, pages 551-556, XP002059197</p> <p>see the whole document</p> <p>-----</p>	1-5,9,10

INTERNATIONAL SEARCH REPORT

In international application No.
PCT/US 97/21723

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☒ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

Claims Nos.: 1,6,9-13

because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

In view of the large number of compounds which are defined by formula III (claim 6) and by the generic terms ("capsaicin agonist" and "capsaicin antagonist") of claim 1, the search had to be restricted for economic reasons. The search was limited to the compounds specifically exemplified and/or claimed and to the general idea underlying the application. (see PCT Guidelines, Chapter III, paragraph 2.3).

Remark : Although claims 9-13 are directed to a method of treatment of the human/animal body , the search has been carried out and based on the alleged effects of the compound/composition.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 97/21723

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9004389 A	03-05-90	US 4939149 A AU 619930 B AU 4624189 A CA 2001194 A,C DE 68910834 D DE 68910834 T EP 0439549 A JP 7029919 B JP 3504387 T US 5290816 A	03-07-90 06-02-92 14-05-90 24-04-90 23-12-93 17-03-94 07-08-91 05-04-95 26-09-91 01-03-94
WO 9717077 A	15-05-97	FR 2740682 A	09-05-97
WO 9015055 A	13-12-90	US 5021450 A AT 141585 T AU 5561490 A CA 2057917 A DE 69028203 D DE 69028203 T EP 0474672 A ES 2093645 T JP 9165333 A JP 2694479 B JP 4503360 T	04-06-91 15-09-96 07-01-91 01-12-90 26-09-96 06-03-97 18-03-92 01-01-97 24-06-97 24-12-97 18-06-92
EP 68590 A	05-01-83	US 4460602 A CA 1191862 A	17-07-84 13-08-85